GENOTYPING PROTOCOL
MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu
530-754-MMRRC

Protocol Name: **CR11781 Pdrg1 EXDEL**

Protocol: ***GoTaq® G2 Colorless Master Mix(Promega)***

Reagent/Constituent	Volume (µL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20µM) comF	0.5
Primer 2. (stock concentration is 20µM) wtR	0.5
Primer 3. (stock concentration is 20µM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	**15.00 µL**

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps		Temp (ºC)	Time (m:ss)	# of Cycles
1. Initiation/Melting	HOT START? ☐	94	2:00	**1x**
2. Denaturation		94	0:10	
3. Annealing	steps 2-3-4 cycle in sequence	65 (↓1ºC/cycle)	0:30	**10x**
4. Elongation		68	2:00	
5. Denaturation		94	0:15	
6. Annealing	steps 5-6-7 cycle in sequence	55	0:30	**25x**
7. Elongation		68	2:00 (↑20sec/cycle)	
8. Finish		4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')
1. CR_Pdrg1_comF	GAGTTAGACCAGATGGTAGTGATGG
2. CR_Pdrg1_wtR*	CCAGTCAACCTTGAGAGTGTACC
3. CR_Pdrg1_mutR	GGAACATAAGGGCAGAACTATTGC

Electrophoresis Protocol:

Agarose: **1.5%** **V:** **90**

Estimated Running Time: **90** min.

Primer Combination	Band (bp)	Genotype
1 & 2,1 & 3	323,918	wildtype
1 & 3	302	mutant

Allele Description: Exon 2 ENSMUSE00001305179.2 and flanking splicing regions were constitutively deleted from the Pdrg1 Gene ENSMUSG00000027472 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 616bp deletion from Chr2:152856223-152856436, 152855816-152856217 GRCm39 was sequence verified. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

