

GENOTYPING BY PCR PROTOCOL
MUTANT MOUSE REGIONAL RESOURCE CENTER: UC DAVIS
 2795 2nd Street, Suite 400, Davis, CA 95618
mmrrc@ucdavis.edu
 530-754-MMRRC

NAME OF PCR: Sanger MirKO ES Cell Line Mir187 - LRPCR **MMRRC #** 034444-UCD

Protocol: SequelPrep long PCR kit, Invitrogen A10498

Reagent/ Constituent	Volume (μL)
Sterile H2O	13.24
10X Buffer (green tube)	2
Enhancer A (Red tube)	1
Enhancer B (Yellow tube)	1
DMSO (brown tube)	.2
gene specific primer (10uM)	0.5
universal primer (10uM)	0.5
Enzyme (black tube)	0.36
DNA extracted with <input type="checkbox"/> NaOH <input checked="" type="checkbox"/> Proteinase K <input checked="" type="checkbox"/> Other: Qiagen DNEasy	1
TOTAL VOLUME OF REACTION:	20μL

Comments on protocol:

May decrease Elongation time for shorter PCR fragments per manufacturers suggestion i.e. 1 minute/kb of sequence.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? <input type="checkbox"/>	93	3:00	1
2. Denaturation	93	0:15	} 8x
3. Annealing	68 to 60 (↓1°C/cycle)	0:30	
4. Elongation	68	9:00	
5. Denaturation	93	0:15	} 32x
6. Annealing	60	0:30	
7. Elongation	68	9:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')
1: 5' common rev	atagcatatcattatcacgaagtattcactgg
2: 5' gene-specific (LR1)	tctttattagagaccttgggatataagatt
3: 3' common fwd	tctagaaagtataggaacttccatggtc
4: 3' gene-specific (LR4)	tataaaaatgggctttgagatcagaagtat

Electrophoresis Protocol:

% Agarose: 0.8 V: 90

Estimated Running Time (min): 90

Primer Combination	Band (kb)	Genotype
1 and 2 (5')	5270	Targeted
3 and 4 (3')	3429	Targeted

Animal Genotyping - Designing Primers for Short Range PCR (SRPCR)

- Go to <http://www.knockoutmouse.org/martsearch/search?query=mir+TV> database and search the gene name i.e. Mir100
- In IKMC Targeted Projects—click on the “View Details”
- Under “# Targeted Non-Conditional Clones”—click on the “Genbank file” and copy into a sequence analysis software such as VectorNTI (Invitrogen) or other analysis tool.
- Locate the vector specific fwd primer “tctagaaagtataggaacttccatggc” and design a reverse complement ~ 200-400 bases downstream (within homology arm). Name “3’ gsp rev” and note expected amplicon size.
- Locate the “LR2” region on the sequence file and design a fwd primer ~ 100-200 bp upstream and name “5’ gsp fwd”. This will pair with the 3’ gsp rev and give a reaction on the wildtype (WT) allele only with this protocol and always slightly larger by a few hundred bp compared to the KO reaction.
- To calculate the expected WT amplicon size: Blast this entire sequence against the mouse genome rather in NCBI or Ensembl and retrieve the contig sequence. Input the contig sequence into your DNA sequence analysis software and determine the exact amplicon size of the 3’ gsp rev and 3’ gsp fwd reaction.
- Contact Brandon Willis at bjwillis@ucdavis.edu at anytime for assistance.