

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu
530-754-MMRRC

Protocol Name: CR10632 Vcan iDex

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (µL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20µM) comF	0.5
Primer 2. (stock concentration is 20µM) wtR	0.5
Primer 3. (stock concentration is 20µM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 µL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? <input type="checkbox"/>	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Electrophoresis Protocol:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5%	V: 90
1. CR_Vcan_comF	TTCCAATCCATTCTCACATTCAGCA	Estimated Running Time: 90 min.	
2. CR_Vcan_wtR	GCATCACCTACATCATCGGGATGT	Primer Combination	Band (bp)
3. CR_Vcan_mutR	TCGTGTTTTCCACATGTTCCACATATCA	1 & 2, 1 & 3	346, 559
		1 & 3	333
			mutant

Allele Description: Exon 3 ENSMUSE00000229348 had 226bp deleted from the 94th coding nucleotide of exon 3 through the 319th coding nucleotide from the Vcan gene ENSMUST00000109546.8 using CRISPR Cas9 gene editing technology in mouse zygotes. Subsequent founders were backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals.

*wtR primer untested (ePCR verified)

*CR_Vcan_wtR is most likely not necessary

